

Déviations pour les temps locaux d'auto-intersection.

Programme des Communications.

6-10 Décembre 2010, CIRM Luminy

Mini-courses: 5h

Erwin Bolthausen

Title: On the volume of the intersection of two Wiener sausages

Abstract: The lectures will give an overview of the attempts to derive the exact $x \rightarrow \infty$ behavior of

$$P(|R^{(1)} \cap R^{(2)}| \geq x)$$

where $R^{(1)}, R^{(2)}$ are the ranges of two independent infinite length random walks in dimension $d \geq 5$, and on the similar question for the Wiener sausage. The conjecture is that

$$P(|R^{(1)} \cap R^{(2)}| \geq x) = \exp[-c(d) x^{(d-2)/d}],$$

with a fairly explicitly known constant. Although, there is a fair amount of evidence for the conjecture, there still is no proof for it.

First lecture: The Khanin-Mazel-Sinai-Shlosman result [3] on the intersection of the ranges of two independent random walks. Detailed proof. Introduction into the results of the paper [1]. Outline of the Donser-Varadhan approach to the volume of the Wiener sausage.

Second lecture: Presentation of some of the key arguments in [1]: The large deviation result. The “Swiss cheese picture”. The concentration of measure argument. Derivation of the upper bound.

Third lecture: Discussion of the analytical properties of the variational problem, including a proof of the important leakage of mass for $d \geq 5$. Introduction into the two-path problem analyzed in [2]: The upper bound.

Forth lecture: Continuation of the analysis of the two-path problem. Solution of the analytic variational problem leading to the variational formula for $c(d)$. Discussion of the open infinite time-horizon problem.

References

- [1] van den Berg, M., Bolthausen, E., and den Hollander, F.: *Moderate deviations for the volume of the Wiener sausage*. Ann. Math. **153**, 355-406 (2001)
- [2] van den Berg, M., Bolthausen, E., and den Hollander, F.: *On the volume of the intersection of two Wiener sausages*. Ann. Math. **159**, 741-783 (2004)
- [3] Khanin, K.M., Mazel, A.E., Shlosman, S.B., and Sinai, Ya. G.: *Loop condensation effects in the behavior of random walks*, in “The Dynkin Festschrift” (M. Freidlin, ed.), Progr. Probab. **34**, Birkhäuser, Boston, 1994, pp. 167-184

Xia Chen

Title Large deviations for the intersection local times of Brownian motions and random walks

Abstract: The sample path intersection has long been of interest to physicists and mathematicians. It presents a physically relevant model for real world phenomena such as random polymers and quantum field. On the other hand, its analysis has provided mathematical challenges. Thus studying the behavior of intersection local times and related functionals is both physically relevant and often requires a variety of new mathematical ideas. In this lecture we focus on the large deviation problems arising from this area.

The self-intersection local time

$$Q_n = \#\{(j, k); 1 \leq j < k \leq n \text{ and } S(j) = S(k)\}$$

of the random walk $S(n)$ on \mathbb{Z}^d measures the self-intersection of the random trajectory $\{S(1), \dots, S(n)\}$; while the range defined as

$$R_n = \#\{S(1), \dots, S(n)\}.$$

The self-intersection of a single random walk is closely related to the mutual intersection between two independent and identically distributed random walks $S(n)$ and $\tilde{S}(n)$. The relevant quantities in mutual intersection are mutual intersection local time

$$I_n = \#\{(j, k) \in [1, n]; S(j) = \tilde{S}(k)\}$$

and the range intersection

$$J_n = \#\left(\{S(1), \dots, S(n)\} \cap \{\tilde{S}(1), \dots, \tilde{S}(n)\}\right).$$

The notion of the intersection local times can be extended from random walks to Brownian motions in the “sub-critical dimensions” (which mean different things for different type of intersections). Compared with their discrete counterparts, the Brownian intersection local times need a more technical setup. On the other hand, there are more analytical tools available to deal with continuous case. In the spirit of invariance principle, the study in the Brownian cases turns out to be a crucial step toward the later success in the discrete cases, in addition to the importance for its own sake.

This lecture presents some progresses recently made on the large deviations for sample path intersections and addresses the following topics:

1. Definition and renormalization of Brownian intersection local times.
2. Le Gall’s moment identity and method of high moment asymptotics
3. LDP by Feynman-Kac formula.
4. Large deviations for Q_n , R_n , I_n , J_n and the relations among these results.
5. Unsolved problems.

References

Bolthausen, E. (1999). Large deviations and interacting random walks. *Lecture Notes in Math.* **1781** 1-124.

Chen, X. *Random Walk Intersections: Large Deviations and Related Topics*. Math. Surv. Mono. **157**, Providence 2009.

Le Gall, J-F. (1992). Some properties of planar Brownian motion. *École d'Été de Probabilités de Saint-Flour XX. 1990. Lecture Notes in Math* **1527** 111-235. Springer, Berlin.

Lectures: 45mn

Frank den Hollander

Title: On the collision local time of two transient random walks

Abstract: In Birkner, Greven and den Hollander (2010), a quenched large deviation principle (LDP) was established for the empirical process of words obtained by cutting an i.i.d. sequence of letters into words according to a renewal process. This LDP can be applied to prove that the radius of convergence of the moment generating function of the collision local time of two independent copies of a symmetric and strongly transient random walk on \mathbb{Z}^d with $d \geq 1$, both starting from the origin, strictly increases when we condition on one of the random walks, both in discrete time and in continuous time. The same is expected to hold when the random walk is transient but not strongly transient. The presence of these gaps implies the existence of an intermediate phase for the long-time behaviour of a class of coupled branching processes, interacting diffusions, respectively, directed polymers in random environments.

Michiel van den Berg

Title: Asymptotics of the heat exchange and some conjectures of M. V. Berry

Abstract: Let K be a compact subset in Euclidean space \mathbb{R}^m , and let $E_K(t)$ denote the total amount of heat in $\mathbb{R}^m \setminus K$ at time t , if K is kept at fixed temperature 1 for all $t \geq 0$, and if $\mathbb{R}^m \setminus K$ has initial temperature 0. For two disjoint compact subsets K_1 and K_2 we define the heat exchange $H_{K_1, K_2}(t) = E_{K_1}(t) + E_{K_2}(t) - E_{K_1 \cup K_2}(t)$. We obtain the leading asymptotic behaviour of $H_{K_1, K_2}(t)$ as $t \rightarrow 0$ under mild regularity conditions on K_1 and K_2 . We discuss some conjectures of M. V. Berry and show how the renormalised heat content of a region D in \mathbb{R}^m determines the length of the shortest closed periodic geodesic in D .

Dima Ioffe

Title: Random walks in attractive potentials: The case of critical drifts.

Abstract: We consider random walks in attractive potentials - sub-additive functions of their local times. An application of a drift to such random walks

leads to a phase transition: If the drift is small than the walk is still sub-ballistic, whereas the walk is ballistic if the drift is strong enough. The set of sub-critical drifts is convex with non-empty interior and can be described in terms of Lyapunov exponents (Sznitman, Zerner). Recently it was shown that super-critical drifts lead to a limiting speed. We shall explain that in dimensions $d \geq 2$ the transition is always of the first order. (Joint work with Y.Velenik)

Wolfgang König

Title: Upper tails of self-intersection local times of random walks: survey of proof techniques

Abstract: The asymptotics of the probability that the self-intersection local time of a random walk on \mathbb{Z}^d exceeds its expectation by a large amount is a fascinating subject because of its relation to some models from Statistical Mechanics, to large-deviation theory and variational analysis and because of the variety of the effects that can be observed. However, the proof of the upper bound is notoriously difficult and requires various sophisticated techniques. We survey some heuristics and some recently elaborated techniques and results.

Peter Mörters

Title: Intersections of random walks in supercritical dimensions

Abstract: In high dimensions two independent simple random walks have only a finite number of intersections. I will present a joint paper with Xia Chen, in which we determine the exact upper tail behaviour of the intersection local time.

Yueyun Hu

Title: Aldous' conjecture on a killed branching random walk.

Abstract: This talk is based on a joint work with Elie Aidekon and Olivier Zindy. Consider a branching random walk on the real line with an killing barrier at zero: starting from a nonnegative point, particles reproduce and move independently, but are killed when they touch the negative half-line. The population of the killed branching random walk dies out almost surely in both critical and subcritical cases, where by subcritical case we mean that the rightmost particle of the branching random walk without killing has a negative speed and by critical case when this speed is zero. We investigate

the total progeny of the killed branching random walk and give its precise tail distribution both in the critical and subcritical cases, which solves an open problem of D.Aldous.

Balint Toth

Title: Superdiffusive bounds on random walks and diffusions with long memory in the critical dimension

Abstract: We prove superdiffusivity with multiplicative logarithmic corrections for a class of models of random walks and diffusions with long memory. The family of models includes the true (or myopic) self-avoiding random walk, self-repelling Durrett-Rogers polymer model and diffusion in the curl-field of (mollified) massless free Gaussian field in 2D. We adapt methods developed in the context of bulk diffusion of ASEP by Landim-Quastel-Salmhofer-Yau (2004). (This is joint work in progress with Benedek Valkó (U Wisconsin, Madison))

Senya Shlosman

Title: Ising model fog drip: the shallow puddle – $o(N)$

Abstract: This joint work with Dima Ioffe is a sequel to our paper Ising model fog drip: the first two droplets (2008).

We study the model of the stalagmite formation. It is a result of the dew-fall effect, when the concentration of the vapor exceeds the saturation point. It turns out that the growth process has discontinuities, when a new atomic monolayer is created spontaneously. An interesting feature of the process is that the size of each newly born monolayer has to exceed some critical size $C_{cr}N$, where N is the linear size of our 3D system. The study boils down to the investigation of the ensemble of the nested random loops in 2D, which are under the influence of two competing mechanisms: entropic repulsion and weak attraction.

Yvan Velenik

Title: Percolation with a line of defects

Abstract: We consider the Bernoulli bond percolation process on the d -dimensional lattice, with occupation probability p for all edges, except those along one line (say, the first coordinate axis), for which the occupation probability is p' .

We assume that p is below the critical value, $p < p_c(d)$. We first prove that the probability that two vertices x and y along the line are connected decays exponentially fast in $|y - x|$ for any value of $p' < 1$. Denote by $\xi_{p,p'}$ the corresponding rate of exponential decay and write $\xi_p = \xi_{p,p}$. Since $\xi_{p,p'} = \xi_p$, for all $p' \leq p$, we can define $p'_c(p, d) = \sup\{p' : \xi_{p,p'} = \xi_p\}$.

We prove that, in dimensions 1, 2 and 3, $p'_c = p$, while in higher dimensions $p < p'_c < 1$. We also analyze the behavior of $\xi_p - \xi_{p,p'}$ as p' decreases toward p in dimensions 1, 2 and 3.

Finally, we derive sharp asymptotics of pure exponential form for the probability of connecting two distant vertices along the line, valid for all $p < p_c$ and all $p' > p'_c$.

Results of the type above are well-known for effective models, in which the central object (here the cluster connecting two distant vertices along the line) is replaced by the path of a random walk and the pinning potential by a suitable exponential functional of the local time at the origin. As far as we know, our results constitute the first analysis of such problems beyond the framework of effective models (apart from the exact computations in the 2D Ising model that triggered the study of these effective models).

Based on joint work with Sacha Friedli and Dmitry Ioffe.

Talks of Wednesday Morning

Anne-Laure Basdevant

Title: Vertex reinforced jump processes on Galton-Watson trees.

Abstract: A vertex reinforced jump process on a tree is a continuous time random walk which jumps to a neighbouring vertex with rate proportional to the local time at that vertex plus a constant. In this talk, we give a criterion to determine whether the walk is recurrent or transient depending on the parameters of the model.

Hubert Lacoin

Title: Superdiffusivity for directed polymer in correlated random environ-

ment

Abstract: The directed polymer in random environment models the behavior of a polymer chain in a solution with impurities. It is a particular case of random walk in random environment. In $1 + 1$ dimensional environment it has been shown by Petermann that this random walk is superdiffusive. We show superdiffusivity properties are reinforced where there are long ranged correlation in the environment and that super diffusivity also occurs in higher dimensions.

Achim Klenke

Title: Multiple Non-Intersection Exponents of Planar Brownian Motion

Abstract: Consider the paths of three planar Brownian motions killed when they leave the unit circle. It is well known that all three paths intersect (i.e., they produce triple points) if all motions are started at the origin. What is the probability p_R that the paths do *not* intersect when started at random positions at the boundary of a small circle with radius R ?

It is shown that

$$p_R \approx R^\xi \quad \text{as } R \downarrow 0,$$

where ξ is the so-called non-intersection exponent. The exact numerical value is unknown and is estimated via a Monte-Carlo simulation. For a slightly more general setup with different numbers of Brownian motions, we obtain rigorous inequalities, precise results and/or approximate values via simulations.

The situation is quite different (in terms of the quality of results) from the one with double points. Here, the corresponding exponents for the non-intersection of m Brownian motions of one type with n Brownian motions of the other type are known exactly due to work of Lawler, Werner and Schramm. (Joint work with Peter Mörters.)

Alexandre Gaudillière

Title: Fluctuation for internal DLA

Abstract: We show logarithmic and sub-logarithmic fluctuations with respect to a spherical asymptotic shape for internal DLA in dimension two and larger than or equal to three respectively.

Marc Wouts

Title: A first order localization/delocalization transition in charged poly-

mers

Abstract: We study a quenched charged-polymer model, introduced by Garel and Orland in 1988, that reproduces the folding/unfolding transition of biopolymers. We prove that, below the critical inverse temperature, the polymer is delocalized in the sense that: (1) The rescaled trajectory of the polymer converges to the Brownian path; and (2) The partition function remains bounded. At the critical inverse temperature, we show that the maximum time spent at points jumps discontinuously from 0 to a positive fraction of the number of monomers, in the limit as the number of monomers tends to infinity. Finally, when the critical inverse temperature is large, we prove that the polymer collapses in the sense that a large fraction of its monomers live on four adjacent positions, and its diameter grows only logarithmically with the number of the monomers. Our methods also provide some insight into the annealed phase transition and at the transition due to a pulling force; both phase transitions are shown to be discontinuous.

Nikos Zygouras

Title: Strong Disorder in Semi-directed Random Polymers.

Abstract : Semi-directed, random polymers can be modeled by a simple random walk on Z^d in a random potential $-(\lambda + \beta\omega(x))_{x \in Z^d}$, where $\lambda > 0$, $\beta > 0$ and $(\omega(x))_{x \in Z^d}$ is a collection of i.i.d., nonnegative random variables. We identify situations where the annealed and quenched costs, that the polymer pays to perform long crossings are different. In these situations we show that the polymer exhibits localization.